

*Spring 2017*

# CSCI 621: **Digital Geometry Processing**

## 9.1 **Surface Parameterization**





Hao Li

<http://cs621.hao-li.com>

# Modeling



# Modeling



# Viewpaint

The creation of a 3D assets surface, including that surface's color, texture, opacity, and reflectivity (or specularly).


# Viewpoint

Rango: Creating creature scale textures in ZBrush...




# Viewpaint




(Wrinkle Pass)



# Color Maps





# Wet Maps






# bump Maps



# Motivation


## Texture Mapping



Levy et al.: *Least squares conformal maps for automatic texture atlas generation*, SIGGRAPH 2002.

# Motivation


## Normal Mapping




# Motivation




Texture Mapping



Normal Mapping




Detail Transfer



Morphing



Mesh Completion




Editing



Databases




Remeshing






Surface Fitting

# Motivation




# Motivation




# Mesh Parameterization

Find a 1-to-1 mapping between given surface mesh and 2D parameter domain






# Unfolding Earth






# Spherical Coordinates




$$\begin{bmatrix} \theta \\ \phi \end{bmatrix} \mapsto \begin{bmatrix} \sin \theta \sin \phi \\ \cos \theta \sin \phi \\ \cos \phi \end{bmatrix}$$

# Desirable Properties

Low distortion



Bijjective mapping



# Cartography



orthographic



stereographic

↑  
preserves angles  
= conformal



Mercator



Lambert

↑  
preserves area  
= equiareal

Floater, Hormann: *Surface Parameterization: A Tutorial and Survey*,  
Advances in Multiresolution for Geometric Modeling, 2005

# More Maps



Mollweide-Projektion



Mercator-Projektion




Zylinderprojektion nach Miller




Hammer-Aitoff-Projektion




Peters-Projektion




Längentreue Azimuthalprojektion



Stereographische Projektion




Behrmann-Projektion




Senkrechte Umgebungsperspektive



Robinson-Projektion



Hotise Oblique Mercator-Projektion



Sinusoidale Projektion



Gnomonische Projektion



Flächentreue Kegelprojektion





Transverse Mercator-Projektion



Cassini-Soldner-Projektion


# Demo: Parameterization



# Recall: Differential Geometry

## Parametric surface representation

$$\mathbf{x} : \Omega \subset \mathbb{R}^2 \rightarrow \mathcal{S} \subset \mathbb{R}^3$$
$$(u, v) \mapsto \begin{pmatrix} x(u, v) \\ y(u, v) \\ z(u, v) \end{pmatrix}$$



## Regular if

- Coordinate functions  $x, y, z$  are smooth
- Tangents are linearly independent


$$\mathbf{x}_u \times \mathbf{x}_v \neq \mathbf{0}$$

# Definitions

**A regular parameterization  $\mathbf{x} : \Omega \rightarrow \mathcal{S}$  is**

- **Conformal** (angle preserving), if the angle of every pair of intersecting curves on  $\mathcal{S}$  is the same as that of the corresponding pre-images in  $\Omega$ .
- **Equiareal** (area preserving) if every part of  $\Omega$  is mapped onto a part of  $\mathcal{S}$  with the same area
- **Isometric** (length preserving), if the length of any arc on  $\mathcal{S}$  is the same as that of its pre-image in  $\Omega$ .

# Distortion Analysis



**Jacobian transforms infinitesimal vectors**

$$d\mathbf{x} = \mathbf{J}d\mathbf{u} \quad \mathbf{J} = \begin{pmatrix} x_u & x_v \\ y_u & y_v \\ z_u & z_v \end{pmatrix}$$

$$\|d\mathbf{x}\|^2 = (d\mathbf{u})^T \mathbf{J}^T \mathbf{J} d\mathbf{u} = (d\mathbf{u})^T \mathbf{I} d\mathbf{u}$$



# First Fundamental Form

Characterizes the surface locally

$$\mathbf{I} = \begin{pmatrix} \mathbf{x}_u^T \mathbf{x}_u & \mathbf{x}_u^T \mathbf{x}_v \\ \mathbf{x}_u^T \mathbf{x}_v & \mathbf{x}_v^T \mathbf{x}_v \end{pmatrix}$$

Allows to measure on the surface

- Angles  $\cos \theta = (\mathbf{du}_1^T \mathbf{I} \mathbf{du}_2) / (\|\mathbf{du}_1\| \cdot \|\mathbf{du}_2\|)$
- Length  $ds^2 = \mathbf{du}^T \mathbf{I} \mathbf{du}$
- Area  $dA = \det(\mathbf{I}) du dv$


# Isometric Maps

**A regular parameterization  $\mathbf{x}(u, v)$  is isometric, iff its first fundamental form is the identity:**

$$\mathbf{I}(u, v) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

**A surface has an isometric parameterization iff it has zero Gaussian curvature**


# Cylinder




# Conformal Maps (A-Similar-AP)

A regular parameterization  $\mathbf{x}(u, v)$  is conformal, iff its first fundamental form is a scalar multiple of the identity:

$$\mathbf{I}(u, v) = s(u, v) \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$




# Conformal Flow



# Equiareal Maps

A regular parameterization  $\mathbf{x}(u, v)$  is equiareal, iff the determinant of its first fundamental form is 1:

$$\det(\mathbf{I}(u, v)) = 1$$



# Relationships

**An isometric parameterization is conformal and equiareal, and vice versa:**

**isometric  $\Leftrightarrow$  conformal + equiareal**

**Isometric is ideal, but rare. In practice, people try to compute:**


- Conformal
- Equiareal
- Some balance between the two

# Harmonic Maps

- A regular parameterization  $\mathbf{x}(u, v)$  is harmonic, iff it satisfies

$$\Delta \mathbf{x}(u, v) = 0$$

- isometric  $\Rightarrow$  conformal  $\Rightarrow$  harmonic
- Easier to compute than conformal, but does not preserve angles





# Harmonic Maps

- A harmonic map minimizes the Dirichlet energy

$$\int_{\Omega} \|\nabla \mathbf{x}\|^2 = \int_{\Omega} \|\mathbf{x}_u\|^2 + \|\mathbf{x}_v\|^2 \, du \, dv$$


- Variational calculus then tells us that

$$\Delta \mathbf{x}(u, v) = 0$$



- If  $\mathbf{x} : \Omega \rightarrow S$  is harmonic and maps the boundary  $\partial\Omega$  of a convex region  $\Omega \subset \mathbb{R}^2$  homeomorphically onto the boundary  $\partial S$ , then  $\mathbf{x}$  is one-to-one.

# Parameterization Goal

- Piecewise linear mapping of a discrete 3D triangle mesh onto a planar 2D polygon
- Slightly different situation: Given a 3D mesh, compute the inverse parameterization



# Floater's Parameterization



# Floater's Parameterization

- For Quadrilateral Patch
- Fix the parameters of the boundary vertices on a unit square
- Derive the bijection  $\mathbf{u}$  for each of the interior vertices  $\mathbf{v}_i$  by solving

$$u(\mathbf{v}_i) = \sum_{k \in \mathcal{V}(i)} \lambda_{i,k} u(\mathbf{v}_k)$$

where  $\lambda_{i,k}$  satisfies shape preserving criteria

$$\text{and } \sum_{k \in \mathcal{V}(i)} \lambda_{i,k} = 1, \quad i = 1, 2, \dots, n$$

# Floater's Algorithm

- Compute for each  $i$  the  $\lambda_{i,k}, k \in v(i)$
- Compute a local parameterization for  $v(i)$  that preserves the aspect ratio of the angle and length
- Compute  $\lambda_{i,k}, k \in v(i)$  that satisfies

Shape preserving criteria

$$\text{and } \sum_{k \in v(i)} \lambda_{i,k} = 1, \quad i = 1, 2, \dots, n$$

- Solve the sparse equation for  $u(v_i), i = 1 \dots n$

$$u(v_i) = \sum_{k \in v(i)} \lambda_{i,k} u(v_k)$$

# Discrete Harmonic Maps

- Map the boundary  $\partial S$  homeomorphically to some (convex) polygon  $\partial\Omega$  in the parameter plane
- Minimize the Dirichlet energy of  $\mathbf{u}$  by solving the corresponding Euler-Lagrange PDE

$$\Delta_S \mathbf{u} = 0$$


- Requires discretization of Laplace-Beltrami
- Compare to surface fairing

# Discrete Harmonic Maps

- System of linear equations

$$\forall v_i \in \mathcal{S} : \sum_{v_j \in \mathcal{N}_1(v_i)} w_{ij} (\mathbf{u}(v_j) - \mathbf{u}(v_i))$$

$$w_{ij} = \cot \alpha_{ij} + \cot \beta_{ij}$$



- Properties of system matrix:
  - Symmetric + positive definite  $\rightarrow$  unique solution
  - Sparse  $\rightarrow$  efficient solvers

# Discrete Harmonic Maps

- But...
  - Does the same theory hold for discrete harmonic maps as for harmonic maps?
  - In other words, is it possible for triangles to flip or become degenerate?



# Convex Combination Maps

- If the linear equations are satisfied

$$\sum_{v_j \in \mathcal{N}_1(v_i)} w_{ij} (\mathbf{u}(v_j) - \mathbf{u}(v_i))$$

and if the weights satisfy

$$w_{ij} > 0 \quad \wedge \quad \sum_{v_j \in \mathcal{N}_1(v_i)} w_{ij} = 1$$

then we get a convex combination mapping.

# Convex Combination Maps

- Each  $\mathbf{u}(v_i)$  is a convex combination of  $\mathbf{u}(v_j)$

$$\mathbf{u}(v_i) = \sum_{v_j \in \mathcal{N}_1(v_i)} w_{ij} \mathbf{u}(v_j)$$

- If  $\mathbf{u} : \mathcal{S} \rightarrow \Omega$  is a convex combination map that maps the boundary  $\partial\mathcal{S}$  homeomorphically to the boundary  $\partial\Omega$  of a convex region  $\Omega \subset \mathbb{R}^2$ , then  $\mathbf{u}$  is one-to-one.

# Convex Combination Maps

- Uniform barycentric weights

$$w_{ij} = 1/\text{valence}(v_i)$$


- Cotangent weights ( $> 0$  if  $\alpha_{ij} + \beta_{ij} < \pi$ )

$$w_{ij} = \cot(\alpha_{ij}) + \cot(\beta_{ij})$$

- Mean value weights


$$w_{ij} = \frac{\tan(\delta_{ij}/2) + \tan(\gamma_{ij}/2)}{\|\mathbf{p}_j - \mathbf{p}_i\|}$$

(no negative weights, even for obtuse angles)




# Convex Combination Maps


- Comparison




original  
mesh



uniform  
weights



cotan  
weights  
(shape preserving)




mean  
value

# Fixing the Boundary


- Choose a simple convex shape
  - Triangle, square, circle
- Distribute points on boundary
  - Use chord length parameterization

Fixed boundary can create high distortion







# Open Boundary Mappings

- Include boundary vertices in the optimization
- Produces mappings with lower distortion




# Open Boundary Mappings




# Need disk-like topology

- Introduce cuts on the mesh






# Naive Cut, Numerical Problems




# Smart Cut, Free Boundary



# Texture Atlas Generation

- Split model into number of patches (atlas)
  - because higher genus models cannot be mapped onto plane and/or
  - because distortion, the number of patches will be too high eventually



Levy, Petitjean, Ray, Maillot: *Least Squares Conformal Maps for Automatic Texture Atlas Generation*, SIGGRAPH, 2002





# Texture Atlas Generation

- Split model into number of patches (atlas)
  - because higher genus models cannot be mapped onto plane and/or
  - because distortion, the number of patches will be too high eventually



Levy, Petitjean, Ray, Maillot: *Least Squares Conformal Maps for Automatic Texture Atlas Generation*, SIGGRAPH, 2002

# Non-Planar Domains




(a) [Alexa 2000]

(b) [Haker et al., 2000]

(c) [Isenburg et al., 2001]

seamless, continuous parameterization of genus-0 surfaces

# Global Parameterization – Range Images



# Constrained Parameterizations




Levy: *Constraint Texture Mapping*, SIGGRAPH 2001.

# Literature

- Book, Chapter 5
- Hormann et al.: Mesh Parameterization, Theory and Practice, Siggraph 2007 Course Notes
- Floater and Hormann: Surface Parameterization: a tutorial and survey, advances in multiresolution for geometric modeling, Springer 2005
- Hormann, Polthier, and Sheffer, Mesh Parameterization: Theory and Practice, SIGGRAPH Asia 2008 Course Notes



# Next Time



Decimation

<http://cs621.hao-li.com>

# Thanks!

